

Correlation between the orbital positions of Pluto/ Neptune and solar activity in the past 8650 years

Sacha P. Dobler, 2021 updated 6.4. 2022

Abstract:

Yearly orbital positions of the four Jovian planets (the gas giants) and Pluto were compared to solar activity proxies: to sunspot numbers (SSN) for the past 8650 years and to radio carbon calibration data of the past 12,000 years. Correlation coefficients were calculated between the angular distance of each planet (or accumulated angular distances of several planets) from 12 different orbital positions and the solar activity of the corresponding year.

Weak but very consistent negative correlations were found between the angular distance of Pluto as well as the accumulated angular distances of Neptune and Pluto from the region of 265° ecliptic longitude and the 40-year moving average of Sunspot Number: sunspot number increases when either Pluto or both Pluto and Neptune are closer to the 265° position (one degree from the orientation of the galactic center) and is lowest when they are closer to the 85° position, with a gradual transition between these points. The relationship is strongest for Pluto observed independently: $r = -0.09022$, $p < 10^{-16}$.

Smaller correlations are found for the accumulated angular distances of Pluto and Neptune from the 265° position, and correlation coefficients below - 0.03 ($p < 0.01$) were obtained for Neptune when observed independently. No other planet or combination of two or more planets yielded statistically significant correlations $r > 0.003$.

Possible contributing interactions of the sun and the planets with charged particles and magnetic fields near the orientation of the galactic center and solar system bow shock are discussed.

Introduction

Previous research in the relationship between solar- and geomagnetic activity with solar system object alignments/ geometry has usually included the planets, with focus on the four gas giants. Existing papers deal with the gravitational effects involving the planets' position in relation to one another and the sun and barycentric perturbations affecting the sun, whereas the planets' positions in relation to the solar system have not been investigated.

In this paper, Pluto was observed with the four gas giants and the positions of the planets were not only analyzed in relation to one another, but in their angular distances from all orientations in the solar system.

Background

Various models have been proposed involving planetary position geometry and solar activity. One of the presented models involve the cycles of the four giant planets in the Solar System: Jupiter, Saturn, Uranus and Neptune, another only Uranus and Neptune, yet another the alignment of Earth, Venus, and Jupiter.

- Early theoretical arguments for the planets as significant outer agents to the Sun go back to the 1970s. These models *"are supported by observations showing that planetary tides follow a pattern correlating with the solar cycle in the last three and a half centuries"*.¹

- Mörner 2012 asserted that Solar variability in emission of luminosity and Solar Wind is driven by a planetary beat, because the Solar-Planetary system behaves like a multi-body system constantly adjusting their motions around a common center of gravity. *"The solar influence on terrestrial climate and environments seems primarily to go via the interaction of the solar wind with the Earth's magnetosphere."*²

- McCracken et al found: Low cosmic-ray intensity (higher solar activity) occurred when Uranus and Neptune were in superior conjunction (mutual cancellation), while high intensities occurred when Uranus–Neptune were in inferior conjunction (additive effects). *"Many of the prominent peaks in the PCR Fourier spectrum can be explained in terms of the Jose cycle, and the occurrence of barycentric anomalies."*³

- During the period 1874 – 2017, the Morlet wavelet power spectrum of the north–south asymmetry of sunspot-group area and that of the mean absolute difference of the orbital positions of **the 4 giant planets** are found to be similar.⁴

- Geoff J. Sharp proposed the Angular Momentum perturbation and modulation is a direct product of the **outer gas giants (Uranus & Neptune)**. The authors accurately predicted in 2013 solar cycle 24 [peaked in 2014] & 25 [started 2020] will be heavily reduced in sunspot activity resembling a similar pattern to solar cycles 5 & 6 during the Dalton Minimum (1790-1830).⁵ Solar Cycle 24 has already turned out to be the lowest in a century and has been considered as a precursor of a new extended solar minimum. ⁶ Lockwood, 2013 *estimated solar activity is "falling faster than at any time in the last 9300 years."*⁷

- A team of researchers from the HZDR found that the tidal forces [of the Synchronized Solar Dynamo] are strongest when **Earth, Venus, and Jupiter** align, and that this alignment occurs every 11.07 years - falling at the same time as the solar minimum. ⁸ They traced back solar cycles from between the years 1000 and 2009 CE, and found an impressively strong link between solar activity and the movements of the planets in that time.

- Andreu et al, 2012 found an excellent agreement between the long-term cycles in proxies of solar activity and the periodicities in the planetary torque and also that some periodicities remain phase-locked over 9400 years. Based on these observations, the authors proposed that the long-term solar magnetic activity is modulated by planetary effects. ⁹ *"Since no successful physical mechanism was suggested to explain these correlations, the possible link between planetary motion and solar activity has been largely ignored."*

Method:

Planetary positions for the past 12,000 years were generated by the Institute of Applied Astronomy of the Russian Academy of Sciences (IAA RAS) using ephemerides EPM2017H (-10106-3036):EPM2017 and EPM2017h: ¹⁰ The IAA ephemerides are concurrent with the NASA JPL DE431 with deviations of up to 1 degree per 10,000 years.

Data from the JPL Horizon online calculator were not included, as they deviate substantially from the JPL DE431 data. SSN values and JPL DE431 data of the orbital positions of Pluto yielded a correlation of 0.26 for the past 450 years, whereas the JPL horizon data yield a correlation of 0.29 for the same time span. ¹¹

Reconstructed sunspot numbers for the years 6755BC to 1894AD were obtained from Wu et al, 2018. ¹² ¹³ The reconstructed SSN and ¹⁴C production are negatively correlated at $r = -0.56$.

Detrended ¹⁴C calibration curves (Reimer et al, 2013) were also compared with planetary positions over the past 12,000 years. Further, Southern Hemisphere ¹⁴C calibration data from McCormac FG, et al 2014 were analyzed. ¹⁴

For planets and their positions that yielded correlation coefficients $r > 0.003$, the accumulated angular distances for each combination of planets from each reference point in the ecliptic were compared to the above proxies of solar activity. All orbital positions are given in °ecliptic longitude.

Results:

Sunspot Number (40-year moving average of SSN) is overall highest when Pluto is closest to the 265° orientation in the Solar System (from 6755BC to 1894AD). Smaller correlations are found for the accumulated angular distances of Pluto and Neptune from this position ($r = -0.8480$; $p < 10^{-14}$), and lower correlation coefficients of $r = -0.03025$ ($p = 0.004$) were obtained for Neptune when observed independently. In this paper, the 265° orientation (one degree from the orientation of the galactic center) was dubbed the *Front End* of the Solar System while the 85° orientation was dubbed the *Back End*. The relationship gradually increases with increasing proximity to the 265° position, decreases to below $r = -0.004$, (not statistically significant, $p > 0.6$) at the 180° and 90° positions and is reversed in the opposite direction, i.e. solar activity is lowest when Pluto and Neptune are closer to the 85° position ($r = 0.08480$; $p < 10^{-14}$), or when Pluto is closest to 85° ($r = 0.09022$; $p < 10^{-16}$).

The correlation of the angular distance of Pluto from the 265° orientation, when observed independently, and SSN is: $r = -0.09022$, $p < 10^{-16}$ for the past 8650 years (r square = 0.008139). No other planets or combination of two or more planets yielded statistically significant correlations with either SSN or any of the other proxies of solar activity of $r > 0.003$.

Radio Carbon production:

When detrended ^{14}C calibration data (Reimer et al, 2013¹⁵, and McCormac et al 2014¹⁶) were compared with planetary positions over the past 12,000 years, the position of the four giant planets or Pluto showed no statistically significant relationships with Radio Carbon production.

The fluctuations in atmospheric ^{14}C are caused by three major factors:

1) changes in the intensity of the geo- magnetic dipole moment, 2) heliomagnetic modulation, i.e., modulation by the magnetic field imbedded in the solar wind, and 3) changes in the carbon cycle. A prescient article by Lingenfelter and Ramaty (1970) also included the production of ^{14}C by solar flares and supernova. “¹⁷




Figure 1 Correlation coefficients of Sunspot Number (40-year moving average) and angular distance from the orbital positions of the same year for: NE =Neptune; PL= Pluto, NE+PL= added angular distances of the two bodies.

Conclusion:

The Planets' (and Pluto's) positions in respect to the heliosphere may be used to predict long-term fluctuations in the amplitude of solar cycles and especially the non-periodical grand solar cycles. Further investigations on the effects of Pluto and Neptune on solar- and geomagnetic activity may help to achieve better predictability not only of solar output, but of the associated changes in climate and social mood.

The Pluto/Neptune - solar activity connection may contribute to the understanding of physical processes in solar system models involving electromagnetic exchanges between the sun, the outer heliosphere and the interstellar wind, possibly involving planets as conductors or capacitors. The termination shock is twice as far from the sun as Pluto and Neptune and the region within is traditionally believed to be affected only in an insignificant way by the interstellar medium and magnetic field.

However, a cloud of entering interstellar dust with particle size of $0.0316 \mu\text{m}$ concentrates at the termination shock centering around 260° , reaching well into the orbits of Neptune and Pluto, with much higher densities than the ambient interstellar dust outside the heliopause (Slavin et al 2012, see fig. 2).¹⁸ The grains tend to be positively charged by the solar UV radiation once they get within ~ 100 AU of the sun.

Discussion:

If future research should find the relationship to have a causal aspect, it is highly probable that Pluto's mass is not sufficient to influence solar activity by gravitational mechanisms/ perturbations of the solar system barycenter. The observation of Neptune's position independently from Pluto yielded considerably lower correlations as compared to Pluto, even though they also point towards the *Front End*. Neptune and Pluto are tidally locked in a 2:3 synchronization and never come closer than 2600

the heliosphere has been measured by at least 11 different spacecrafts as of 2013.³⁰ The flow, coming from Earth's perspective from the constellation Scorpius, "has probably changed direction by several degrees since the 1970s".³¹

Further, a broad belt of energetic protons with a nonthermal pressure comparable to that of the local interstellar magnetic field, is centered at approximately 260 degrees ecliptic longitude extending from north to south and looping back through approximately 80 degrees. "The belt appears to be ordered by the local interstellar magnetic field."³²

W. I. Axford and S. T. Suess, 1994 stated; "It is natural to assume that the relative velocity [of the Sun and the heliosphere] might be parallel to the motion of the Sun with respect to the nearer stars, namely from the solar apex (right ascension ~ 271 degrees, declination ~ 30 degrees) with a speed of ~ 20 km/s. However, typical interstellar clouds have random (line of sight) velocities of ~ 10 km/s or more and the direction could easily be 30 degrees or more from the solar apex - which turns out to be the case."³³

The solar wind pressure - a combined measure of its speed and density - increased by approximately 50 percent in 2014 (at the peak of the last solar maximum) and remained that way for several years thereafter.³⁴

Voyager I is travelling in the almost exact direction (longitude) against the interstellar wind (the direction of the Front end) but at a steeper ascension in respect to the solar system plain.

The spacecraft crossed the heliopause, or the edge of the heliosphere, in August 2012. Heading in a different direction, Voyager 2 crossed another part of the heliopause in November 2018.³⁵

The heliosphere itself is thought to reside in the Local Interstellar Cloud inside the Local Bubble, which is a region in the Orion Arm of the Milky Way Galaxy.

Outside the heliosphere there is a forty-fold increase in plasma density.³⁶

The local interstellar medium (LISM), which is partly ionized plasma, interacts with the solar wind (SW) plasma and forms the heliospheric interface. At the heliospheric interface, solar wind and interstellar plasma, interplanetary and interstellar magnetic fields, interstellar atoms, galactic and anomalous cosmic rays (GCRs and ACRs) and pickup ions play prominent roles."³⁷

- Geoff J. Sharp 2013, who proposed the Angular Momentum perturbation and modulation of solar activity is a direct product of the **outer gas giants (Uranus & Neptune)**, accurately predicted in 2013 solar cycle 24 [peaked in 2014] & 25 [started 2020] will be heavily reduced in sunspot activity resembling a similar pattern to solar cycles 5 & 6 during the Dalton Minimum (1790-1830).³⁸ SC 24 has already turned out to be the lowest in a century and has been considered as a precursor of a new extended solar minimum.³⁹ Lockwood, 2013 *estimated solar activity is "falling faster than at any time in the last 9300 years."*⁴⁰

Sacha P. Dobler 2021; contact sacha.dobler@gmx.net




Figure 3: Inverted accumulated angular distances of Pluto and Neptune for the 265 ° position and SSN (40-year moving average) 6755BC to 1894AD (dates labeled as year before 1894AD).

Notes:

¹ Grandpierre, Attila, 1996: On the origin of solar cycle periodicity; *Astrophysics and Space Science* volume 243, pages 393–400 (1996)

² Mörner, Nils-Axel 2012: PLANETARY BEAT, SOLAR WIND AND TERRESTRIAL CLIMATE; *Solar Wind: Emission, Technologies and Impacts* – p. 47-66, 2012; *Paleogeophysics and Geodynamics*, Stockholm, Sweden

³ McCracken K. G. Steinhilber 2014: Evidence for Planetary Forcing of the Cosmic Ray Intensity and Solar Activity Throughout the Past 9400 Years; *Coronal Magnetometry* pp 585-607

⁴ Javaraiah J. 2020: Long-term Periodicities in North–south Asymmetry of Solar Activity and Alignments of the Giant Planets; *Solar Physics* volume 295, Article number: 8 (2020)

⁵ Sharp, G. J. 2013: Are Uranus & Neptune Responsible for Solar Grand Minima and Solar Cycle Modulation?; *International Journal of Astronomy and Astrophysics*; Vol.3 No.3 (2013),

⁶ Zharkova, V.V. 2015: Heartbeat of the Sun from Principal Component Analysis and prediction of solar activity on a millennium timescale. *Scientific Report*, 5 *Scientific Reports* volume 5

⁷ Newscientist, Fred Pearce, 2013: *Solar activity heads for lowest low in four centuries*; 1 November 2013 By <https://www.newscientist.com/article/dn24512-solar-activity-heads-for-lowest-low-in-four-centuries/>

⁸ Stefani F. et al, 2019: A Model of a Tidally Synchronized Solar Dynamo; *Solar Physics* volume 294, Article number: 60 (2019)

⁹ Abreu J. A et al 2012: Is there a planetary influence on solar activity? *A&A* 548, A88 (2012)

¹⁰ <http://iaaras.ru/en/dept/ephemeris/epm/2017/>

¹¹ <ftp://ssd.jpl.nasa.gov/pub/eph/planets/>

¹² Wu, C. J., Usoskin I. G., Krivova, N., et al. Solar activity over nine millennia: A consistent multi-proxy reconstruction; *A&A* 615, A93 (2018); <https://doi.org/10.1051/0004-6361/201731892>; ESO 2018

¹³ Sunspot group numbers timeSerie (865 rows)

<https://vizier.u-strasbg.fr/viz-bin/VizieR?-source=J/A+A/615/A93>

¹⁴ McCormac, F., Hogg, A., Blackwell, P., Buck, C., Higham, T., & Reimer, P. (2004). SHCal04 Southern Hemisphere Calibration, 0–11.0 Cal Kyr BP. *Radiocarbon*, 46(3), 1087-1092. doi:10.1017/S0033822200033014

¹⁵ M, Reimer RW, Richards DA, Scott EM, Southon JR, Turney CSM, van der Plicht J. 2013: IntCal13 and MARINE13 radiocarbon age calibration curves 0-50000 years calBP; *Radiocarbon* 55(4). DOI: 10.2458/azu_js_rc.55.16947

¹⁶ McCormac FG, et al 2014: SHCal04 Southern Hemisphere Calibration 0 - 11.0 cal kyr BP; *Radiocarbon* 46, 1087-1092

¹⁷ Damon, Paul et al 2000: RADIOCARBON CALIBRATION AND APPLICATION TO GEOPHYSICS, SOLAR PHYSICS, AND ASTROPHYSICS; *RADIOCARBON*, Vol 42, Nr 1, 2000, p 137–150 © 2000 by the Arizona Board of Regents on behalf of the University of Arizona

¹⁸ Slavin Jonathan D. et al 2012: Trajectories and Distribution of Interstellar Dust Grains in the Heliosphere; *The Astrophysical Journal*, 760:46 (15pp), 2012 November 20 doi:10.1088/0004-637X/760/1/46 © 2012.

¹⁹ NASA History; SP-345 Evolution of the Solar System; 8.5.1. Neptune-Pluto; 01/25/2021 <https://history.nasa.gov/SP-345/ch8.htm>

²⁰ Benson G. S. et al 1971: Resonances in the Neptune-Pluto System

April 1971 *The Astronomical Journal* 76(2); DOI:10.1086/111100; SourceNTRS

²¹ https://www.vanderbilt.edu/AnS/physics/astrocourses/ast201/pluto_tombaugh.html

the discovery of Pluto; vanderbilt.edu; *Astronomy 201: The Solar System*; Spring 2003

²² Jonathan D. Slavin et al 2012: Trajectories and Distribution of Interstellar Dust Grains in the Heliosphere; *The Astrophysical Journal*, 760:46 (15pp), 2012 November 20 doi:10.1088/0004-637X/760/1/46 © 2012.

²³ Liewer P. C. et al., 2000: *NASA's interstellar probe mission*, *Space Technology and Applications International Forum-2000*, edited by M. S. El-Genk (AIP, Maryland, 2000), p. 911.

²⁴ Reid, M.J.; Brunthaler, A. (2004). "The Proper Motion of Sagittarius A*^{*}". *The Astrophysical Journal*. 616 (2): 874, 883.

²⁵ : <https://www.vercalendario.info/en/how/convert-ra-degrees-hours.html>

²⁶ NASA fact sheet PLUTO: <https://nssdc.gsfc.nasa.gov/planetary/factsheet/plutofact.html>

²⁷ Opher M. Stone E. C & Gombosi T. I: The Orientation of the Local Interstellar Magnetic Field <https://arxiv.org/pdf/0705.1841.pdf>

²⁸ Lallement, R., et al, 2005, *Deflection of the interstellar neutral hydrogen flow across the heliospheric interface*, *Science* 307, 1447–1449

²⁹ Ben-Jaffel, L et al: . 2012: *Uncovering the magnetic environment of our solar system*; *A&A*; Volume 546, October 2012 A78; DOI <https://doi.org/10.1051/0004-6361/201219637>

³⁰ NASA 2013: Eleven Spacecraft Show Interstellar Wind Changed Direction Over 40 Years; NASA Official: Brian Dunbar <https://www.nasa.gov/content/goddard/interstellar-wind-changed-direction-over-40-years>

³¹ Zell, Holly (6 March 2015). "Interstellar Wind Changed Direction Over 40 Years".

³² Krimigis SM, Mitchell DG, Roelof EC, Hsieh KC, McComas DJ. Imaging the interaction of the heliosphere with the interstellar medium from Saturn with Cassini. *Science*. 2009 Nov 13;326(5955):971-3. doi: 10.1126/science.1181079. Epub 2009 Oct 15. Erratum in: *Science*. 2010 Apr 9;328(5975):170. PMID: 19833914.

³³ Axford; W. I. Suess. S. T. 1994; *Spacecraft to explore the outer heliosphere* Volume 75, Issue 50; 13 December 1994; Pages 587-594

³⁴ NASA Jun 6, 2018; nasasibexobs; As Solar Wind Blows, Our Heliosphere Balloons; Rob Garner
<https://www.nasa.gov/feature/goddard/2018/as-solar-wind-blows-our-heliosphere-balloons>

³⁵ Jet Propulsion Laboratory; Two Interstellar Travelers; Dec 10, 2018; <https://www.jpl.nasa.gov/images/two-interstellar-travelers>

³⁶ NASA Spacecraft Embarks on Historic Journey Into Interstellar Space; Greicius, Tony (5 May 2015)
https://www.nasa.gov/mission_pages/voyager/voyager20130912.html

³⁷ Izmodenov, V.V. 2001: The Outer Heliosphere: The Next Frontiers; in COSPAR Colloquia Series, 2001;
<https://www.sciencedirect.com/topics/physics-and-astronomy/interstellar-plasma>

³⁸ Sharp, G. J. 2013: Are Uranus & Neptune Responsible for Solar Grand Minima and Solar Cycle Modulation?;
International Journal of Astronomy and Astrophysics; Vol.3 No.3 (2013),

³⁹ Zharkova, V.V. 2015: Heartbeat of the Sun from Principal Component Analysis and prediction of solar activity
on a millennium timescale. Scientific Report, 5 Scientific Reports volume 5

⁴⁰ Newscientist, Fred Pearce, 2013: *Solar activity heads for lowest low in four centuries*; 1 November 2013
By <https://www.newscientist.com/article/dn24512-solar-activity-heads-for-lowest-low-in-four-centuries/>